

The anamorphic universe

This content has been downloaded from IOPscience. Please scroll down to see the full text.

JCAP10(2015)001

(<http://iopscience.iop.org/1475-7516/2015/10/001>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 128.112.66.66

This content was downloaded on 24/11/2015 at 15:27

Please note that [terms and conditions apply](#).

The anamorphic universe

Anna Ijjas^{a,1} and Paul J. Steinhardt^{a,b}

^aPrinceton Center for Theoretical Science, Princeton University,
Princeton, NJ, 08544 U.S.A.

^bDepartment of Physics, Princeton University,
Princeton, NJ, 08544 U.S.A.

E-mail: aijjas@princeton.edu, steinh@princeton.edu

Received July 19, 2015

Revised August 4, 2015

Accepted August 9, 2015

Published October 1, 2015

Abstract. We introduce “anamorphic” cosmology, an approach for explaining the smoothness and flatness of the universe on large scales and the generation of a nearly scale-invariant spectrum of adiabatic density perturbations. The defining feature is a smoothing phase that acts like a contracting universe based on some Weyl frame-invariant criteria and an expanding universe based on other frame-invariant criteria. An advantage of the contracting aspects is that it is possible to avoid the multiverse and measure problems that arise in inflationary models. Unlike ekpyrotic models, anamorphic models can be constructed using only a single field and can generate a nearly scale-invariant spectrum of tensor perturbations. Anamorphic models also differ from pre-big bang and matter bounce models that do not explain the smoothness. We present some examples of cosmological models that incorporate an anamorphic smoothing phase.

Keywords: modified gravity, alternatives to inflation, physics of the early universe

ArXiv ePrint: [1507.03875](https://arxiv.org/abs/1507.03875)

¹Corresponding author.

Contents

1	Introduction	1
2	Essentials of anamorphic cosmology	3
3	Advantage over previous models	5
4	Conditions for an anamorphic smoothing phase	5
5	Scalar-tensor formulation	9
6	Perturbations	10
6.1	Adiabatic and tensor modes from the anamorphic phase	10
6.2	No multiverse	12
7	Examples	13
7.1	Approximate solutions for $\epsilon(N)$	13
7.2	$V_J(\phi) = V_0 e^{B\phi}$	14
7.3	$V_J(\phi) = V_0 e^{B\phi - C\phi^p}$	14
8	Completing the cosmological scenario	16
8.1	Simple (one-time) anamorphosis	17
8.2	Cyclic anamorphosis	20
9	Discussion	23
A	Cosmological background in Jordan and Einstein frames	24
A.1	Jordan-frame representation	24
A.2	Einstein-frame representation	25
A.3	Matter contribution and contraction	26
B	Conditions for successful anamorphosis	26

1 Introduction

Observations show the large-scale structure of the universe to be flat, smooth and scale-free [24] and the spectrum of primordial curvature perturbations to be nearly scale-invariant, adiabatic and Gaussian. Inflationary cosmology attempts to explain these large-scale properties by invoking a period of rapid accelerated expansion, but this approach leads to an initial conditions problem and a multiverse problem that each make the theory unpredictable. Ekpyrotic cosmology invokes a period of slow contraction that makes the universe smooth and flat while avoiding initial conditions and multiverse problems, but it requires an entropic mechanism using two or more scalar fields to generate the nearly scale-invariant spectrum of curvature perturbations.

In this paper, we introduce “anamorphic cosmology” that combines elements and advantages of both earlier approaches. Based on some invariant criteria, the cosmological

background during the smoothing phase behaves like a contracting universe that homogenizes, isotropizes and flattens without introducing an initial conditions or multiverse problem; and, based on other invariant criteria, the cosmological background behaves like an expanding universe that can directly generate a scale-invariant spectrum of super-horizon adiabatic perturbations using a single scalar field. Because the impression of the cosmological background depends on the perspective (i.e., the Weyl frame), we refer to models of this type as “anamorphic.”

An anamorphic cosmology integrates a combination of concepts described in the forthcoming sections:

- massive particles whose mass m in any Weyl frame has a different time-dependence than the Planck mass M_{Pl} , i.e., m/M_{Pl} is time varying (section 2);
- two Weyl-invariant quantities, Θ_m and Θ_{Pl} , that characterize the contraction or expansion relative to the characteristic matter scale (e.g., Compton wavelength) or to the gravitational (Planck) scale, respectively (section 2);
- suppression of the anisotropy due to time-varying m/M_{Pl} (section 4);
- a scalar-tensor theory realization of the anamorphic phase (section 5);
- a kinetic coupling that causes Θ_m and Θ_{Pl} to have opposite signs (section 5);
- generation of nearly scale-invariant adiabatic, Gaussian *density* perturbations in an anamorphic contracting phase governed by a single scalar field (section 6);
- generation of nearly scale-invariant *gravitational-wave* perturbations in an anamorphic contracting phase (section 6);
- a novel, anamorphic mechanism for averting a multiverse (section 6.2);
- two kinds of bounces (termed Θ_m -bounce and Θ_{Pl} -bounce) in which Θ_m or Θ_{Pl} reverse sign from negative to positive (section 8);
- transition from contraction to expansion and vice versa without violation of the null-energy condition (NEC) (section 8.1);
- or, in some cases, non-singular bounce from contraction to expansion with ghost-free, stable NEC violation (section 8.2);
- incorporation of the elements above into a geodesically complete cyclic cosmology in which the anamorphic phase, matter-radiation creation, and dark-energy domination repeat at regular intervals (section 8.2).

Although some of these elements have been considered by other authors (most notably by Piao [36], Wetterich [44], Li [30], and Domenech and Sasaki [13], as we will discuss), the novelty here is how they work together with the other elements to form a complete cosmological scenario.

This paper is organized as follows. Sections 2–6 describe the anamorphic phase during which the universe is smoothed and adiabatic curvature perturbations are generated. In sections 2–3, we describe the essential components of anamorphic cosmology, the advantages compared to other cosmological scenarios, and the cosmological background. Section 4 details

the conditions that the background solution must satisfy to smooth and flatten the universe and produce squeezed quantum states. Then, in section 5, we present a simple scalar-tensor field theory that realizes the anamorphic conditions. In this paper, we focus on cases for which there exists a frame in which the anamorphic scalar field ϕ couples only to itself and the Ricci scalar; generalizations of anamorphic cosmology will be discussed in forthcoming publications. The second-order action describing the perturbations and the conditions for obtaining a nearly scale-invariant spectrum are given in section 6. The analysis shows how anamorphosis circumvents an earlier no-go theorem [12] stating that adiabatic modes always decay during contracting phases if mediated by a single scalar field. We then explain how the multiverse is avoided. We conclude with examples that demonstrate how to incorporate an anamorphic smoothing phase into a complete cosmological scenario. A simple model with a single anamorphic phase is analyzed in section 8.1. In section 8.2, we describe the construction of a model in which the smoothing phase, the bounce and the transition to a hot expanding phase repeat at regular intervals in a geodesically complete, cyclic cosmology.

2 Essentials of anamorphic cosmology

Anamorphic cosmology relies on having two essential components during the smoothing phase: time-varying masses and a combination of Weyl-invariant signatures that incorporate aspects of contracting and expanding backgrounds.

The first component is that particle masses and/or the Planck mass have different time dependence in any Weyl frame. For simplicity, we will consider models in which matter-radiation consists of massive dust. The frame-invariant action for a single particle is

$$S_p = \int m ds, \quad (2.1)$$

where ds is the line element and m may vary with time. For a homogeneous and isotropic universe, we can assume a Friedmann-Robertson-Walker (FRW) metric $ds^2 = -dt^2 + a^2(t)dx_i dx^i$ where t is the FRW time, $a(t)$ is the scale factor and the Hubble parameter is $H \equiv d \ln a / dt$.

To distinguish cosmological models, it is convenient to introduce the two dimensionless Weyl-invariant quantities:

$$\Theta_m \equiv \left(H + \frac{\dot{m}}{m} \right) M_{\text{Pl}}^{-1}, \quad (2.2)$$

$$\Theta_{\text{Pl}} \equiv \left(H + \frac{\dot{M}_{\text{Pl}}}{M_{\text{Pl}}} \right) M_{\text{Pl}}^{-1}. \quad (2.3)$$

Intuitively, Θ_m measures the physical expansion or contraction of the cosmological background as measured relative to a ruler (or any object made of matter); in the cases considered in this paper, Θ_m also measures the congruence of geodesics. Θ_{Pl} measures the evolution relative to the Planck mass, which is important for analyzing the generation of scalar and tensor metric perturbations.

In standard cosmology with constant particle mass m and Planck mass M_{Pl} , the two invariants are equal, $\Theta_m M_{\text{Pl}} = \Theta_{\text{Pl}} M_{\text{Pl}} = H$, so no distinction is made. In particular, both invariants are positive during an inflationary smoothing phase and negative during an ekpyrotic smoothing phase. One can introduce a Weyl transformation of the metric in either case to re-express the action such that M_{Pl} or m or both are time-dependent. In fact, as

	Anamorphosis	Inflation	Ekpyrosis
Θ_m	< 0	> 0	< 0
background	contracts	expands	contracts
Θ_{Pl}	> 0	> 0	< 0
curvature perturbations	grow	grow	decay

Table 1. Distinguishing the anamorphic, inflationary, and ekpyrotic scenarios: in conventional inflationary and ekpyrotic models, both the particle mass m and the Planck mass M_{Pl} are constant, and the two Weyl-invariants Θ_m and Θ_{Pl} are both equal to the Hubble parameter (in Einstein frame). Consequently, both invariants are positive during an inflationary smoothing phase and both are negative during an ekpyrotic smoothing phase. In anamorphic models, Θ_m , which characterizes the behavior of the cosmological background, is negative, as in ekpyrotic cosmology; and Θ_{Pl} , which characterizes the generation and evolution of scalar and tensor metric perturbations, is positive, as in inflationary models.

Wetterich [44] and Li [30] have recently emphasized, it is possible to transform expanding models to a frame where the Hubble parameter H is negative. However, the advantage of the variables Θ_m and Θ_{Pl} is that they are frame invariant (resolving an issue also discussed in [13]). For example, Wetterich’s formulation of the action can still be unambiguously recognized as a version of expansion during all phases by computing these Weyl-invariant quantities and showing them both to be positive. Analogous constructions are possible in ekpyrotic models, in which case both invariants are negative during the smoothing phase.

The second essential component of an anamorphic phase is that Θ_m is negative (as in ekpyrotic models) and Θ_{Pl} is positive (as in inflationary models), which is possible in principle if m and M_{Pl} are time-dependent, as can be seen from eqs. (2.2)–(2.3). In section 4, we will discuss how to construct models whose equation of state during the anamorphic phase leads to the pair of Weyl-invariant quantities having this behavior. Because Θ_m is negative, the cosmological background is physically contracting such that smoothing occurs without creating a multiverse or incurring an initial conditions problem, as in ekpyrotic models. Because Θ_{Pl} is positive, the second-order action describing the generation and evolution of curvature perturbations during the smoothing phase is similar to the case of inflation. Consequently, a nearly scale-invariant spectrum of adiabatic density and gravitational wave perturbations can be generated in models with a single scalar field.

From the invariants, therefore, it is clear that anamorphic models are unambiguously distinctive and, at the same time, share the best traits of inflationary and ekpyrotic cosmology while avoiding the worst. The essential differences are summarized in table 1.

Table 1 suggests a fourth possibility with Θ_m positive and Θ_{Pl} negative. In fact, such a set-up is achievable for different choices of the equation of state and time-varying masses. However, for the purposes of constructing useful cosmological models, this possibility is unpromising since it has the cosmological background behavior of inflation, thereby typically producing a multiverse, and the generation of curvature perturbations, governed by Θ_{Pl} , is made more complicated.

3 Advantage over previous models

An anamorphic cosmology has several advantages over the classic inflationary paradigm [2, 21, 31]. First, unlike inflation [20, 35], initial conditions do not have to be finely-tuned for the smoothing phase to begin. Second, inflation is geodesically incomplete [6], but cyclic anamorphic models can be constructed that are geodesically complete. Third, eternal smoothing [38, 41] is avoided, so there is no multiverse. Hence, the predictions in a given anamorphic model are unambiguous and testable.

Among cosmological models that feature periods of contraction, simple pre-big bang [17–19, 40] and matter bounce models [16, 42] are unstable to anisotropy perturbations and, hence, do not explain smoothness and flatness. Anamorphic and ekpyrotic models do, but they employ different smoothing mechanisms. Ekpyrosis [25] uses a period of ultra-slow contraction in which the pressure, p , exceeds the energy density, ρ ($w \equiv p/\rho > 1$). The anamorphic model relies instead on time-varying masses to suppress the anisotropy and imposes different conditions on the equation of state, as described in the forthcoming sections.

A disadvantage of ekpyrosis is that it is not possible to generate scale-invariant curvature perturbations with a single scalar field during the slow contraction phase [12]. A second scalar field is needed such that one linear combination of the two fields creates scale-invariant entropic perturbations that are converted to curvature perturbations prior to or during the bounce [7, 15, 27, 34]. By contrast, the anamorphic mechanism requires only a single scalar field to do the smoothing, flattening and generation of curvature perturbations. Entropic perturbations and a conversion from entropic to curvature perturbations are not needed.

In the ekpyrotic picture, there is no known way of generating a scale-invariant spectrum of tensor fluctuations. The *primordial* tensor-to-scalar ratio is effectively $r = 0$, a firm prediction that can be proven wrong in future experiments. (N.B. Secondary perturbations are automatically generated in this and all cosmological models by density fluctuations entering the horizon after matter domination; these contribute an effective $r \lesssim 10^{-6}$ [5].) The anamorphic model generates a nearly scale-invariant spectrum of gravitational wave perturbations. Future observations will determine if this is an advantage.

4 Conditions for an anamorphic smoothing phase

As described in section 2, the cosmological background during an anamorphic smoothing phase is described by two Weyl-invariant quantities — $\Theta_m < 0$ and $\Theta_{\text{Pl}} > 0$ — where the first measures contraction with respect to a physical ruler composed of matter and the second measures the expansion with respect to the Planck length, respectively. These quantities can be expressed in terms of two additional frame-invariants:

$$\alpha_m \equiv a \frac{m}{M_{\text{Pl}}^0}, \quad (4.1)$$

$$\alpha_{\text{Pl}} \equiv a \frac{M_{\text{Pl}}}{M_{\text{Pl}}^0}, \quad (4.2)$$

where a is the FRW scale factor, m is the particle mass, M_{Pl} is the reduced Planck mass expressed in a given frame and M_{Pl}^0 is the value of the reduced Planck mass in the frame where it is time independent, e.g., Einstein frame in scalar-tensor theories. The invariant

quantities α_m and α_{Pl} satisfy the relations:

$$\Theta_m = M_{\text{Pl}}^{-1} \frac{\dot{\alpha}_m}{\alpha_m} \quad \text{and} \quad (4.3)$$

$$\Theta_{\text{Pl}} = M_{\text{Pl}}^{-1} \frac{\dot{\alpha}_{\text{Pl}}}{\alpha_{\text{Pl}}}, \quad (4.4)$$

so that $\alpha_{m,\text{Pl}}$ can be viewed as the effective Weyl-invariant scale factors associated with the Weyl-invariant Hubble-like parameters $\Theta_{m,\text{Pl}}$.

During the anamorphic phase, it is most useful to express the first Friedmann equation in a frame-invariant form using Θ_m and α_m :

$$3\Theta_m^2 \left(1 - \frac{d \ln(m/M_{\text{Pl}})}{d \ln \alpha_m}\right)^2 = \frac{\rho_{\Lambda}}{M_{\text{Pl}}^4} + \frac{\rho_m}{M_{\text{Pl}}^4} - \left(\frac{m}{M_{\text{Pl}}}\right)^2 \frac{\kappa}{\alpha_m^2} + \left(\frac{m}{M_{\text{Pl}}}\right)^6 \frac{\sigma^2}{\alpha_m^6}. \quad (4.5)$$

The Friedmann equation describes the contributions of different forms of energy density and curvature to the contraction or expansion rate. The first contribution, $\rho_{\Lambda}/M_{\text{Pl}}^4$, is due to the anamorphic energy density that dominates during the smoothing phase. In the example given in the next section, ρ_{Λ} is ascribed to a non-minimally coupled scalar field ϕ with interaction potential $V_J(\phi)$. The second contribution, $\rho_{\text{matter}}/M_{\text{Pl}}^4$, is due to the matter-radiation energy density in which the matter consists of particles with mass m . The last two contributions are due to the spatial curvature, where $\kappa = (+1, 0, -1)$, and to the anisotropy, parameterized by σ^2 .

The anamorphic combination of $\Theta_m < 0 < \Theta_{\text{Pl}}$ requires that m and/or M_{Pl} be time-dependent such that the invariant mass ratio m/M_{Pl} decreases with time. Note that the factors of m/M_{Pl} in the Friedmann equation suppress the spatial curvature and, even more so, the anisotropy. This is a key feature because suppressing the anisotropy in a contracting universe is essential for avoiding chaotic mixmaster behavior and maintaining homogeneity and isotropy. At some point after the anamorphic phase, to reach consistency with all current cosmological observations and tests of general relativity, the universe must reheat to a high temperature and enter a hot expanding phase in which both m and M_{Pl} become constant, with $M_{\text{Pl}} = M_{\text{Pl}}^0$, the current value of the reduced Planck mass. Throughout this paper, we will use reduced Planck units with $M_{\text{Pl}}^0 \equiv 1$, except where specified otherwise.

Of course, the Friedmann equation during the anamorphic phase can also be described using the invariants Θ_{Pl} and α_{Pl} , as we will do at the end of this section. However, these variables are less useful because they obscure the role of the time-varying masses and the conditions needed after the anamorphic phase to join onto a hot expanding phase.

Next, we consider the generic conditions that Θ_m and α_m must satisfy to resolve the homogeneity and isotropy problems and to generate nearly scale-invariant perturbations.

Smoothing condition. To quantify the condition for smoothing in a model-independent way, we define the effective equation-of-state parameter, ϵ_m , and a mass-variation index q according to the following relations:

$$\epsilon_m \equiv -\frac{1}{2} \frac{d \ln \Theta_m^2}{d \ln \alpha_m}, \quad (4.6)$$

and

$$q \equiv \frac{d \ln(m/M_{\text{Pl}})}{d \ln \alpha_m}. \quad (4.7)$$

Substituting the second relation into eqs. (4.1)–(4.2), we find

$$\frac{d\alpha_{\text{Pl}}}{d\alpha_m} = 1 - q. \quad (4.8)$$

Recalling that, α_m is decreasing and α_{Pl} is increasing in an anamorphic phase, we see that the mass-variation index must satisfy $q > 1$. Note that this constraint implies the condition that the invariant mass ratio m/M_{Pl} is decreasing during the anamorphic phase.

In general, ϵ_m and q can vary with time; however, for simple models, smoothing for an extended period of time is achieved by having them change very slowly during the anamorphic phase; hence, treating them as roughly constant is a useful approximation deep in the anamorphic phase. In the case of nearly constant q , the expression for Θ_m takes a particularly simple form, namely $\Theta_m^2 \propto 1/\alpha_m^{2\epsilon_m}$.

In analogy with ekpyrotic and inflationary models, the conditions for smoothing and flattening an anamorphic universe can be expressed as constraints on the equation-of-state ϵ_m . They require that the anamorphic energy density ρ_A/M_{Pl}^4 dominates all other contributions on the right hand side of the Friedmann equation for an extended amount of time. ‘‘Domination’’ means $\Theta_m^2(1 - q)^2 \propto \rho_A/M_{\text{Pl}}^4$ or, from eq. (4.6),

$$\frac{d \ln (\rho_A/M_{\text{Pl}}^4)}{d \ln \alpha_m} = -2\epsilon_m + \frac{d \ln (1 - q)^2}{d \ln \alpha_m}. \quad (4.9)$$

Note that the second expression on the right hand side of eq. (4.9) is very small, in general, so ρ_A/M_{Pl}^4 is roughly proportional to $1/\alpha_m^{2\epsilon_m}$. In order for eq. (4.9) to hold as α_m shrinks during the anamorphic contracting phase, $2\epsilon_m$ must exceed the corresponding exponents for the spatial curvature and anisotropy terms in the Friedmann equation if they are expressed as powers of $1/\alpha_m^{2\epsilon_m}$. This condition yields a pair of constraints on ϵ_m :

$$\epsilon_m \gtrsim 1 - q \quad \& \quad \epsilon_m \gtrsim 3(1 - q). \quad (4.10)$$

Here, for simplicity, we have neglected the weak time-dependence of q (like the second term on the right hand side of eq. (4.9)). Since $q > 1$, both conditions are satisfied if the first inequality is satisfied, i.e., if $\epsilon_m \gtrsim 1 - q$. (Note that an ekpyrotic contracting phase corresponds to both α_m and α_{Pl} decreasing and constant m/M_{Pl} . Hence, $q = 0$ and the smoothing constraint in eq. (4.10) reduces to $\epsilon_m = \epsilon \equiv (3/2)(1 + w) > 3$, the known result.)

Squeezing of quantum curvature perturbations. To obtain a nearly scale-invariant spectrum of super-horizon curvature perturbations, the cosmological background must have the property that modes whose wavelengths are inside the horizon at the beginning of the smoothing phase can have wavelengths larger than the horizon size by the end of the smoothing phase. The ‘horizon’ is roughly the Hubble radius $|H|^{-1}$ in standard ekpyrotic and inflationary cosmology, and the squeezing condition (modes exiting the horizon scale during smoothing) is that $|aH|$ is increasing.

More generally, the ‘horizon’ is a dynamical length scale that separates smaller wavelengths for which the curvature modes are oscillatory from the large wavelengths for which the curvature modes become frozen. In anamorphic models, the evolution of scalar and tensor metric perturbations are described entirely by the gravitational and scalar field parts of the effective action and do not depend on particle mass, as shown in section 6. Hence, the dynamical length scale is set by Θ_{Pl}^{-1} and the corresponding squeezing condition is that $\alpha_{\text{Pl}}\Theta_{\text{Pl}}$ be increasing,

$$\frac{d|\alpha_{\text{Pl}}\Theta_{\text{Pl}}|}{dt} M_{\text{Pl}}^{-1} > 0, \quad (4.11)$$

which reduces to the standard condition in inflationary and ekpyrotic models. As shown in appendix B, if combined with the second Friedmann equation, the squeezing constraint reduces to the same condition as the smoothing constraint described above in eq. (4.10). Hence, squeezing imposes no additional constraint.

Sufficient anamorphosis. Finally, the anamorphic smoothing phase must last sufficiently long to solve the horizon and flatness problems and generate a broad enough band of scalar and tensor metric perturbations that spans the observable horizon today. Since the metric perturbations are most simply described by α_{P1} and Θ_{P1} , as noted above, it is most straightforward to label modes by the variable N , the number of e -folds of anamorphic phase remaining after the modes become frozen, as measured by the increase in α_{P1} :

$$N \equiv \int_{\alpha_N}^{\alpha_0} d \ln \alpha_{\text{P1}}, \quad (4.12)$$

where $\alpha_N = \alpha_{\text{P1}}(N)$ is the value of α_{P1} at the N th e -fold mark and α_0 is the value at reheating. N runs from some large positive N_{max} to zero during the anamorphic phase. The condition that the duration of the anamorphic phase is sufficient to smooth and flatten the universe is $N_{\text{max}} > 60$.

Other relations. The expansion of the cosmological background relative to fixed rulers (set by the constant particle mass m) is most clearly described by the invariants α_m and Θ_m , as we have done above. However, it is always possible to use the invariants $\Theta_{\text{P1}}, \alpha_{\text{P1}}$, instead. Using these variables simplifies some expressions but has the disadvantage that it obscures the effects of time-varying mass since $\alpha_{\text{P1}} = \alpha_m(M_{\text{P1}}/m)$ absorbs the mass dependence. Also, because α_{P1} is increasing even though the universe is physically contracting, these variables are not useful for determining how to join properly the anamorphic contracting phase to a hot expanding phase. For completeness, though, we repeat here many of the relations above that were expressed in terms of the invariants α_m and Θ_m .

The first Friedmann equation reduces to

$$3\Theta_{\text{P1}}^2 = \frac{\rho_A}{M_{\text{P1}}^4} + \frac{\rho_m}{M_{\text{P1}}^4} - \frac{\kappa}{\alpha_{\text{P1}}^2} + \frac{\sigma^2}{\alpha_{\text{P1}}^6}. \quad (4.13)$$

In analogy with the definition of ϵ_m in eq. (4.6), it is useful to introduce an effective equation-of-state parameter ϵ_{P1} such that

$$\epsilon_{\text{P1}} \equiv -\frac{1}{2} \frac{d \ln \Theta_{\text{P1}}^2}{d \ln \alpha_{\text{P1}}}. \quad (4.14)$$

Using the relation $(1 - q)d\alpha_m = d\alpha_{\text{P1}}$ and the definition of the mass-variation index in eq. (4.7), it is straightforward to show that

$$\epsilon_m - \frac{1}{2} \frac{d \ln(1 - q)^2}{d \ln \alpha_m} = (q - 1)\epsilon_{\text{P1}}, \quad (4.15)$$

such that the smoothing and squeezing constraints reduce to

$$\epsilon_{\text{P1}} < 1. \quad (4.16)$$

In a frame-invariant form, the second Friedmann equation is given as

$$\frac{d\Theta_{\text{P1}}}{dt} M_{\text{P1}}^{-1} = -\frac{1}{2} \frac{\rho_A + p_A}{M_{\text{P1}}^4}, \quad (4.17)$$

where p_A/M_{Pl}^4 is due to the anamorphic pressure. Combining with the first Friedmann equation in eq. (4.5), we find

$$\epsilon_{\text{Pl}} = \frac{3\rho_A + p_A}{2\rho_A}; \quad (4.18)$$

in particular, ϵ_{Pl} is equivalent to the conventional equation of state parameter ϵ .

5 Scalar-tensor formulation

The discussion in previous sections invoked Weyl-invariant variables that describe the contraction or expansion relative to time-varying particle mass and Planck mass. This formulation can be applied to a wide range of models and metrics, as we will explore in future publications. However, to make the ideas concrete, we will henceforth confine our discussion to models in which the anamorphic phase and the transition to a conventional hot expanding phase are mediated by a single scalar field ϕ that is non-minimally coupled to the Ricci scalar R and non-linearly coupled to its kinetic energy density.

The action for our scalar-tensor examples can be expressed as

$$S = \int d^4x \sqrt{-g} \left(\frac{1}{2} M_{\text{Pl}}^2(\phi) R - \frac{1}{2} k(\phi) (\partial_\mu \phi)^2 - V_J(\phi) + \mathcal{L}_m(\phi) \right), \quad (5.1)$$

where $g_{\mu\nu}$ is the metric; R is the Ricci scalar; $k(\phi)$ is the non-linear kinetic coupling function; $V_J(\phi)$ is the potential energy density; and \mathcal{L}_m is a Lagrangian density that describes all other matter and radiation degrees of freedom. Henceforth, we will use the label J to denote frame-dependent quantities in Jordan-frame representation; E to denote Einstein frame; and no designation for quantities that are Weyl-invariant.

In the simple examples considered in this paper, there is a Weyl transformation such that \mathcal{L}_m is independent of ϕ and there is another Weyl transformation that makes M_{Pl} independent of ϕ and \mathcal{L}_m ϕ -dependent. The corresponding equations of motion are given in appendix A. $M_{\text{Pl}}(\phi) \equiv M_{\text{Pl}}^0 \sqrt{f(\phi)}$ is positive definite and approaches the current value $M_{\text{Pl}}^0 \equiv 1$ following the anamorphic phase and through the present epoch. These properties are typical of general scalar-tensor theories. The distinctive feature that enables the anamorphic phase and the transition to and from is the variation of the kinetic coupling, $k(\phi)$, from positive (correct sign) to negative (wrong sign) or vice versa.

The fact that $k(\phi)$ crosses through zero prevents its elimination by a field redefinition. In all cases, the action is constructed so that it is ghost-free. In some scenarios, the action above suffices. In others, we introduce additional Galileon-type terms, $\sim (\partial\phi)^2 \square\phi$, $\sim (\partial\phi)^4$, that play a role before or after the anamorphic phase.

We note that Piao [36] and Li [30] discussed scalar-tensor theories with negative kinetic couplings as a mechanism for generating nearly scale-invariant tensor fluctuations. We will see a similar effect on tensor fluctuations in our anamorphic models. However, Piao's example is explicitly inflationary (in our classification, $\Theta_m = \Theta_{\text{Pl}} > 0$), and Li's is ill-defined since he does not specify the properties of the matter sector, though, he describes his model as dual to inflation. Another difference with Li is that his model maintains the negative kinetic coupling at all stages, including after the reduced Planck mass becomes fixed, which can add some complications after the smoothing phase.)

As we show in appendix B, the conditions for smoothing and flattening the universe and for generating scale-invariant scalar and tensor metric perturbations combined with the

no-ghost condition can be re-expressed in terms of the functions $f(\phi)$, $k(\phi)$ and the equation-of-state parameter ϵ_m or, using eq. (4.15), $\epsilon_{\text{Pl}} \equiv \epsilon$. Combining these constraints, we have

$$0 < 3 + 2k(\phi) \frac{f(\phi)}{(M_{\text{Pl}}^0 f, \phi)^2} < \epsilon < 1. \quad (5.2)$$

The left-hand inequality is the no-ghost condition. As discussed in appendix A, the total kinetic energy including the gravitational contribution (i.e., the kinetic energy in Einstein frame) is given by $(1/2)K(\phi)(\dot{\phi}/M_{\text{Pl}})^2$ with $K(\phi) \equiv (3/2)(M_{\text{Pl}}^0 f, \phi / f)^2 + k/f$. Hence, $K(\phi)$ must be positive definite in order for the field to have the right-sign kinetic term. The middle inequality is the contraction condition ($\Theta_m < 0$); and the third inequality is the smoothing and squeezing constraint ($\epsilon < 1$).

Clearly, given that $f(\phi) > 0$, eq. (5.2) can only be satisfied if the kinetic function

$$k(\phi) < 0 \quad (5.3)$$

during the smoothing phase. As we have emphasized, $k(\phi) < 0$ does not cause any physical instability provided the no-ghost condition, $K(\phi) > 0$, is satisfied, which is already incorporated in eq. (5.2).

Finally, with $d \ln \alpha_{\text{Pl}} = \Theta_{\text{Pl}} (M_{\text{Pl}}/\dot{\phi}) d\phi$ and the expression for ϵ in eq. (B.3), we can rewrite the expression for N in eq. (4.12) as

$$N = \int_{\phi_N}^{\phi_0} \sqrt{\frac{K(\phi)}{2\epsilon}} \frac{d\phi}{M_{\text{Pl}}^0} \gtrsim 60, \quad (5.4)$$

such that the condition of sufficient anamorphosis translates into the constraint that eq. (5.2) holds for $\phi_{60} \leq \phi < \phi_0$. These two conditions on $f(\phi)$, $k(\phi)$ and ϵ are sufficient to have an anamorphic smoothing phase that produces a homogeneous and isotropic cosmological background. As in the case of inflationary and ekpyrotic models, stronger constraints must be imposed in order for the spectrum of perturbations to be nearly scale-invariant, as discussed in the next section.

6 Perturbations

In this section, we discuss the generation of anamorphic scalar and tensor fluctuations and show that adiabatic curvature modes with a (nearly) scale-invariant spectrum are generated. Notably, the generation mechanism requires only a single field, is dynamically stable, results in negligible non-Gaussianity, and produces tensor modes in addition to scalar modes. These are all features that do not occur in ekpyrotic models and have been considered unattainable in bouncing cosmologies [8, 12, 26, 28, 29, 39]. The key reason why they can occur in the anamorphic scenario is that, even though the parameter Θ_m is negative so that the physical background is contracting, Θ_{Pl} is positive. A related effect is that the spectral tilts of both the scalar and tensor perturbation spectra are typically red, whereas one might expect them to be blue in a contracting universe. We will also explain why the anamorphic scenario has no multiverse problem, which is more subtle than the case in ekpyrotic models.

6.1 Adiabatic and tensor modes from the anamorphic phase

The linearized perturbations of our model defined thru the action in eq. (5.1) can be evaluated using the second order actions for both scalar curvature and tensor perturbations.

Curvature perturbations, ζ , are local perturbations in the scale factor that are described by the perturbed metric

$$ds^2 = -dt^2 + a(t)e^{2\zeta(t,x^i)}dx^i dx_i. \quad (6.1)$$

As shown, e.g., in [30], following the standard calculation [3, 32], for the (second-order) action for scalar curvature perturbations ζ we obtain

$$S_{2,S} = \frac{1}{2} \int dt d^3x \frac{\dot{\phi}^2}{\left(H + \dot{M}_{\text{Pl}}/M_{\text{Pl}}\right)^2} K(\phi) M_{\text{Pl}}^2(\phi) a(t) \left[a^2(t) \dot{\zeta}^2 - c_s^2 (\partial_k \zeta)^2 \right], \quad (6.2)$$

where $c_s^2 \equiv 1$ is the speed of sound. Note that, to avoid ghost fluctuations, $K(\phi)$ must be positive definite, which is the same condition that we identified above in eq. (5.2) as the no-ghost constraint. Furthermore, the action for ζ can be put in a frame-independent form:

$$S_{2,S} = \int d\eta d^3x \alpha_{\text{Pl}}^2 \epsilon_{\text{Pl}} \left[(\zeta, \eta)^2 - c_s^2 (\partial_k \zeta)^2 \right], \quad (6.3)$$

where both α_{Pl} and ϵ_{Pl} are defined as in eqs. (2.2) and (4.6) and we have switched to conformal time η , given by $d\eta = dt/a$.

Defining a transverse and traceless tensor perturbation h_{ij} by

$$\delta g_{ij} \equiv a^2 h_{ij}, \quad (6.4)$$

we obtain the analogous quadratic action for tensor perturbations,

$$S_{2,T} = \frac{1}{8} \int d\eta d^3x \alpha_{\text{Pl}}^2 \left[(h_{ij,\eta})^2 - (\partial_k h_{ij})^2 \right]. \quad (6.5)$$

From the definition of the curvature and tensor perturbations and eq. (A.7), it follows that both ζ and h_{ij} are frame invariant as are the prefactors $\alpha_{\text{Pl}}^2, \alpha_{\text{Pl}}^2 \epsilon_{\text{Pl}}$. Hence, the perturbed quadratic actions are invariant under a frame change. As anticipated in earlier sections, although the cosmologically contracting background is most intuitively described in terms of α_m , the perturbations are most intuitively described by α_{Pl} .

In order to obtain nearly scale-invariant spectra of curvature and tensor perturbations, we must choose the couplings, $f(\phi), k(\phi)$ and the potential, $V_J(\phi)$, such that the corresponding pre-factors $\alpha_{\text{Pl}}^2 \epsilon_{\text{Pl}}$ and α_{Pl}^2 scale approximately as η^{-2} , where any small correction to the exponent leads to a tilt.

In an ekpyrotic contracting phase, there is no such option. The pre-factors in the corresponding quadratic actions behave differently: for scalar modes, the pre-factor is proportional to $\alpha_{\text{Pl}}^2 \equiv \alpha_m^2$ (assuming a constant or slowly varying equation of state $\epsilon_{\text{Pl}} \equiv \epsilon_m$) where α_m is very slowly decreasing as the universe contracts. This behavior is clearly different from η^{-2} , which is rapidly growing. This explains why neither tensor modes nor adiabatic scalar modes get amplified during an ekpyrotic phase for models with a single scalar field. With two-fields, there is a mechanism to generate nearly scale-invariant adiabatic perturbations by first generating isocurvature perturbations and then converting, but there is no analogous mechanism for tensor modes so they are predicted to be absent [7, 27].

In the anamorphic contracting phase, the situation is different because the pre-factors depend on $\alpha_{\text{Pl}} \neq \alpha_m$, which is rapidly growing even though the effective scale factor α_m associated with the cosmological background (as measured by Θ_m) is decreasing. In the

limit that ϵ_{Pl} is nearly constant and much smaller than unity, $\alpha_{\text{Pl}}^2 \epsilon_{\text{Pl}}$ and α_{Pl}^2 are nearly proportional to η^{-2} and the spectrum is nearly scale invariant. In section 7, we shall show that it is possible to construct such examples.

We note that the background constraints (smoothing, flattening and squeezing) only require $\epsilon_m > 1 - q$ or $\epsilon_{\text{Pl}} \equiv \epsilon < 1$, but nearly scale-invariant perturbations require stronger conditions, $\epsilon_m \approx 0$ or $\epsilon_{\text{Pl}} \ll 1$ and nearly constant over tens of e -folds of smoothing. The same comment applies to inflation and ekpyrotic models.

It is a common myth that scale-invariance is an automatic outcome of smoothing and flattening. For example, it is sometimes described as a prediction of inflation. In none of these cosmologies is this actually the case. Near scale-invariance, in accordance with observations, imposes significantly tighter constraints on the equation-of-state and its time variation. In the case of inflation, for example, the condition is imposed by requiring that the inflationary phase be nearly de Sitter, which is much stronger than needed to smooth and flatten.

6.2 No multiverse

The multiverse arises in inflation because of the same quantum physics that produces the scale-invariant spectrum described in the previous subsection. Rare quantum fluctuations keep some regions of space in the smoothing phase when a naive estimate would suggest that inflation should have ended. Within instants, the rare regions comprise most of the volume of the universe. The situation repeats eternally, so that only a fractal volume less than three dimensions ever ends inflation. This volume of measure zero containing regions that are no longer inflating is comprised of patches that complete inflation at different times after different random quantum fluctuations have affected the trajectory of the inflaton field that controls the rate of inflation. The different fluctuations lead to different cosmological outcomes. In addition to patches that are flat, smooth, and have nearly scale-invariant perturbation spectra, there are infinitely many patches that are curved, inhomogeneous, anisotropic and with non-scale-invariant fluctuations. By spanning all possible cosmological outcomes, the multiverse effect makes the inflationary scenario unpredictable. There is no generally accepted solution to this problem within inflationary cosmology at present [22].

The multiverse problem does not occur in ekpyrotic cosmology because regions continuing to smooth due to rare fluctuations are contracting rather than expanding. These rare contracting regions shrink away compared to typical regions that complete the contraction phase and transition to an expanding phase.

A similar conclusion holds for anamorphic cosmology, but a more subtle analysis is required. As we have emphasized, during the anamorphic smoothing phase the cosmological background is contracting ($\Theta_m < 0$) as measured by physical rulers comprised of particles with fixed mass m . For rare fluctuations of the scalar field that create patches with negative $\Theta_m < 0$ (within which contraction continues for longer periods compared to typical patches), the situation is the same as in the ekpyrotic case. These rare regions shrink away compared to typical ones. However, as we will illustrate with our simple example in section 8, there can also be rare fluctuations that stop or reverse the evolution of the scalar field and that can flip the sign of Θ_m to positive. Now the rare patch is expanding rather than contracting, which cannot occur in the ekpyrotic picture. We will see that these rare patches are not problematic, though. There is a strong attractor solution that rapidly re-reverses the evolution of the scalar field and the sign of Θ_m before any significant expansion can occur. After the reversal to contraction, the analysis reduces to cases already considered: the rare patches never grow to dominate and there is no multiverse.

7 Examples

In this section, we illustrate some basic properties of anamorphic models using simplistic examples that are only intended to describe the anamorphic contracting phase. In the next section, we describe how to embed such examples in models that describe a complete bouncing cosmology that includes a hot expanding phase following the anamorphic phase.

Let us consider the anamorphic action,

$$S = \int d^4x \sqrt{-g_J} \left(\frac{1}{2} \xi e^{2A\phi} R_J + \frac{1}{2} e^{2A\phi} (\partial_\mu \phi)^2 - V_J(\phi) \right), \quad (7.1)$$

where we set the gravitational coupling $f(\phi) = \xi e^{2A\phi}$ and the kinetic coupling $k(\phi) = -e^{2A\phi}$ with both parameters ξ and A being positive real numbers and we set $M_{\text{Pl}}^0 = 1$. For this family of models, the condition for anamorphic smoothing from eq. (5.2) reduces to the simple form

$$0 < \frac{6A^2\xi - 1}{2\xi A^2} < \epsilon < 1. \quad (7.2)$$

Recall that requiring strict positivity eliminates ghosts despite the fact that $k < 0$.

7.1 Approximate solutions for $\epsilon(N)$

If the smoothing condition in eq. (7.2) is satisfied, we can neglect the contributions from curvature, anisotropy, matter, radiation, etc. and, using the dimensionless “time variable” N defined in eq. (4.12), the equations of motion as given in appendix (A.1–A.3) reduce to

$$\begin{aligned} \frac{\epsilon_{,N}}{\epsilon} &= -2 \frac{\epsilon_{,\phi}}{\sqrt{2\epsilon K(\phi)}} = 2 \frac{3 - \epsilon}{\sqrt{2\epsilon K(\phi)}} \left(\frac{V_{J,\phi}}{V_J} - 2 \frac{f_{,\phi}}{f} + \sqrt{2\epsilon K(\phi)} \right) \\ &= 2(3 - \epsilon) \left(\frac{1}{\sqrt{2\epsilon}} \sqrt{\frac{\xi}{6\xi A^2 - 1}} \left(\frac{V_{J,\phi}}{V_J} - 4A \right) + 1 \right). \end{aligned} \quad (7.3)$$

The solution for ϵ can be approximated analytically if we assume a slowly varying equation of state, i.e.,

$$\frac{\epsilon_{,N}}{\epsilon} \Delta N < 1. \quad (7.4)$$

To satisfy this inequality and the constraint that $\epsilon < 1$ in eq. (7.2), the factor in parentheses in eq. (7.3) must be very small, or equivalently,

$$\epsilon \simeq \frac{1}{2K(\phi)} \left(\frac{d}{d\phi} \ln(V_J(\phi)/f^2(\phi)) \right)^2 = \frac{1}{2} \frac{\xi}{6\xi A^2 - 1} \left(\frac{V_{J,\phi}}{V_J} - 4A \right)^2. \quad (7.5)$$

As noted at the end of section 6.1, a slowly varying equation of state in the anamorphic (or inflationary or ekpyrotic) scenario is not necessary to solve the horizon and flatness problems. For example, it suffices that $\epsilon_{,N} \Delta N < 1 - \epsilon$ in the anamorphic and inflationary models. Rather, it is the empirical constraint that the spectral tilt is observed to differ by only a few per cent from exact scale-invariance that forces the condition that ϵ be so slowly varying and $\ll 1$. A similar statement applies to inflationary and ekpyrotic models.

For simplicity of presentation, we consider a sequence of potentials that can be studied analytically,

$$V_J(\phi) = V_0 e^{B\phi - C\phi^p}, \quad (7.6)$$

for constant V_0 , B and C and integer p , although the results can be easily extended to potentials of the form $V_J \propto e^{g(\phi)}$.

7.2 $V_J(\phi) = V_0 e^{B\phi}$

First, let us consider the case with $C = 0$ and

$$V_J(\phi) = V_0 e^{B\phi}. \quad (7.7)$$

The pre-factor V_0 is a positive real number; it determines the amplitude of scalar perturbations,

$$\delta\rho/\rho(\phi_N) \sim \sqrt{K} \frac{\sqrt{V_J}}{f} \left(\frac{V_{J,\phi}}{V_J} - 2 \frac{f_{,\phi}}{f} \right)^{-1} \sim \sqrt{V_0} \sqrt{\frac{6\xi A^2 - 1}{\xi}} \frac{e^{(1/2)(B-4A)\phi_N}}{\xi|B-4A|}. \quad (7.8)$$

Using eq. (7.3), it is easy to see that the potential in eq. (7.7) corresponds to a constant equation of state,

$$\epsilon \equiv \frac{1}{2} \frac{\xi}{6A^2\xi - 1} (B - 4A)^2. \quad (7.9)$$

The condition for anamorphic contraction in eq. (7.2) reduces to

$$0 < \frac{1}{2}(4A - B)^2 < \frac{6A^2\xi - 1}{\xi} < A|4A - B|, \quad (7.10)$$

where the first inequality is the no-ghost condition ($K(\phi) > 0$), the second inequality is the smoothing condition ($\epsilon < 1$), and the third inequality is the condition for contraction ($\Theta_m < 0$). Note that, in this example, $|4A - B|$ must be strictly greater than zero and non-negligible.

For the purposes of illustration, we introduce a modification that suddenly ends the smoothing phase at $\phi = \phi_0$. (We will present a practical example of this in the next subsection.) Then, the number of e -folds of anamorphic contraction remaining is given by eq. (5.4),

$$N = \frac{6A^2\xi - 1}{\xi} \frac{1}{|4A - B|} \Delta\phi, \quad (7.11)$$

where $\Delta\phi \equiv \phi_0 - \phi_N > 0$. Using ref. [43], the expression for the scalar tilt immediately follows,

$$n_S - 1 = -2\epsilon + \frac{d \ln \epsilon}{dN} = -2\epsilon = \frac{\xi}{6A^2\xi - 1} (B - 4A)^2 = -\frac{|4A - B| \Delta\phi}{N}. \quad (7.12)$$

Observations of the CMB dictate that $n_S - 1 \simeq -1/30$ at $N \simeq 60$, and, thus, for this example, $\Delta\phi \simeq 2/|4A - B|$ and $(6A^2\xi - 1)/\xi \simeq (4A - B)^2(N/2)$. Note that, with this choice of parameters, both the smoothing and the no-ghost constraints in eq. (7.2) are fulfilled. Contraction occurs if, in addition, we require that $|4A - B|(N/2) < A$.

The result demonstrates that, in principle, anamorphic contracting models can generate a nearly scale-invariant spectrum of tensor fluctuations.

7.3 $V_J(\phi) = V_0 e^{B\phi - C\phi^p}$

A simple modification of the previous example is $V_J = V_0 e^{g(\phi)}$ where $g(\phi)$ is positive and increases linearly with ϕ for small ϕ and is negative and decreasing for large ϕ . As a specific example, we consider

$$V_J(\phi) = V_0 e^{B\phi - C\phi^p}, \quad p \gg 2. \quad (7.13)$$

so that the anamorphic phase comes to an end when ϕ grows sufficiently large. The equation of state ϵ begins very small and increases during the anamorphic phase, reaching $\epsilon = 1$ at $\phi = \phi_0 \lesssim 1$ when the anamorphic phase ends.

As in the previous example, the pre-factor V_0 determines the amplitude of scalar perturbations,

$$\delta\rho/\rho(\phi_N) \sim \sqrt{V_0 \frac{6\xi A^2 - 1}{\xi}} \frac{e^{(1/2)(B-4A)\phi_N - (1/2)C\phi_N^p}}{\xi|B - 4A - Cp\phi_N^{p-1}|}. \quad (7.14)$$

The ϕ -dependent equation of state is given by

$$\epsilon \simeq \frac{1}{2} \frac{\xi}{6A^2\xi - 1} (4A - B + Cp\phi^{p-1})^2. \quad (7.15)$$

Note that the equation of state is monotonically growing with increasing $\phi > 0$ if the no-ghost condition, $(6A^2\xi - 1)/\xi > 0$, is satisfied. This ensures that the density-fluctuation spectrum has a red tilt. The anamorphic phase ends at $\epsilon(\phi_0) = 1$, so

$$\phi_0 = \left(\left(\sqrt{2 \frac{6A^2\xi - 1}{\xi}} - 4A + B \right) \frac{1}{Cp} \right)^{1/(p-1)}. \quad (7.16)$$

Note that $\epsilon < 1$ and the smoothing condition is fulfilled for $0 < \phi < \phi_0$.

During the anamorphic phase, there are two regimes of interest: (I) $|4A - B| \gtrsim Cp\phi_0^{p-1} \gg Cp\phi^{p-1}$; and (II) $Cp\phi^{p-1} \gg |4A - B| \simeq 0$:

Case I: if $|4A - B| \gtrsim Cp\phi_0^{p-1} \gg Cp\phi^{p-1}$, the term with coefficient C in the potential in eq. (7.13) is negligible and ϵ is nearly constant except just before the anamorphic phase ends when ϕ is very close to ϕ_0 , where ϕ_0 is given in eq. (7.16). For $\phi_0 \sim 1$ and $N \sim 60$, the prediction for the scalar tilt is

$$n_S - 1 \simeq -|4A - B|/N, \quad (7.17)$$

where the contraction condition can be satisfied by choosing $A > N$. From the second-order action in eq. (6.5), the tensor tilt r for nearly constant $\epsilon_{\text{pl}} \equiv \epsilon$ is given by

$$r \sim -8(n_S - 1), \quad (7.18)$$

which is the same as the standard inflationary consistency relation.

Note that, if the parameters are fit to the current observational constraint on the tilt, $n_S - 1 \sim -1/30$ at $N \approx 60$, then Case I cannot satisfy the current observational bound on $r < 10\%$ [1].

Case II: if $Cp\phi^{p-1} \gg |4A - B| \simeq 0$, the condition for contraction ($\Theta_m < 0$) is given by

$$0 < \frac{6A^2\xi - 1}{\xi} < ACp\phi_N^{p-1}, \quad (7.19)$$

where, in defining the upper bound, we have used the fact that $0 < \phi_N < \phi_0$, i.e., if the physical background contracts for $\phi_N > 0$, it contracts for all $\phi > \phi_N$.

For the number of e -folds of anamorphic contraction, we find

$$N = \frac{6A^2\xi - 1}{\xi} \frac{1}{Cp} \int_{\phi_N}^{\phi_0} \frac{d\phi}{\phi^{p-1}} = \frac{6A^2\xi - 1}{\xi} \frac{1}{Cp} \frac{1}{(p-2)} \left(\frac{1}{\phi_N^{p-2}} - \frac{1}{\phi_0^{p-2}} \right); \quad (7.20)$$

and the expression for the scalar spectral tilt is

$$(n_S - 1)(\phi_N) = -2 \frac{\xi}{6A^2\xi - 1} C p (p-1) \phi_N^{p-2} \left(\frac{C p}{p-1} \phi_N^p + 1 \right). \quad (7.21)$$

Since $(\phi_0/\phi_N)^{p-2} \gg 1$, we have $(n_S - 1)(\phi_N) \simeq -(2/N)(p-1)/(p-2)(1 + C p \phi_N^p/(p-1))$. In order to obtain the observed value of the tilt, $n_S - 1$, p must be large enough that $(p-1)/(p-2) \sim 1$. To satisfy the remaining conditions for successful anamorphosis as stated in eqs. (7.16)–(7.20), it suffices to choose the parameters A, C, ϕ_0 , and ξ as follows,

$$A > \frac{(p-2)N}{\phi_N}; \quad C \ll \frac{1}{\phi_N}; \quad \phi_0 = \left(2 \frac{p-2}{C p} N \right)^{\frac{1}{2(p-1)}} \sqrt{\phi_N}; \quad \frac{6A^2\xi - 1}{\xi} = \frac{1}{2} C^2 p^2 \phi_0^{2(p-1)}. \quad (7.22)$$

Using these parameter constraints, the equation of state and its derivative are given by

$$\epsilon(\phi_N) \simeq \frac{\xi}{2(6\xi A^2 - 1)} C^2 p^2 \phi_N^{2(p-1)} \simeq \left(\frac{\phi_N}{\phi_0} \right)^{2(p-1)} \ll \frac{2}{N} \phi_N^{p-2}, \quad (7.23)$$

$$\frac{d \ln \epsilon}{dN}(\phi_N) \simeq -\frac{2\epsilon_{,\phi}}{\sqrt{2\epsilon} K(\phi)} \simeq -\frac{2}{N}. \quad (7.24)$$

Note that, since $\phi_N \ll 1$ and $p > 2$, ϵ is small compared to its derivative, $|d \ln \epsilon / dN|$. As a result, unlike Case I, it is possible to set the tilt $n_S - 1 \sim d \ln \epsilon / dN$ to the observed value and have the tensor-to-scalar ratio, $r \sim 16\epsilon$ be $\mathcal{O}(1/N^2)$ or smaller, which satisfies the current observational constraint on r .

8 Completing the cosmological scenario

As exemplified in the previous section, the anamorphic phase provides a smoothing mechanism that renders the universe homogeneous, isotropic and spatially flat, generating nearly scale-invariant, super-horizon curvature perturbations. For a complete cosmological scenario, though, we must also explain how the anamorphic phase ends and smoothly connects to standard expanding big-bang evolution. This includes specifying how $f(\phi)$ and $k(\phi)$ must behave during the transition.

During the anamorphic phase, $\Theta_m < 0$ and $\Theta_{P1} > 0$. In the expanding big-bang phase, $\Theta_m = \Theta_{P1} > 0$. To connect the two phases, it is necessary that Θ_m switch sign. We will use the term Θ_m -*bounce* to refer to the moment when Θ_m switches sign from negative to positive; that is, a transition from α_m decreasing to α_m increasing. (A switch from positive to negative Θ_m , which corresponds to α_m switching from increasing to decreasing, will be called a Θ_m -*reversal*. The analogues for Θ_{P1} and α_{P1} will be called Θ_{P1} -*bounce* and Θ_{P1} -*reversal*.) The first model we will describe in section 8.1 is a *simple (one-time) scenario* that (i) begins with an anamorphic smoothing phase; (ii) undergoes a Θ_m -bounce and reheats; and (iii) transitions to standard hot big-bang cosmology with decelerating expansion. As with all current inflationary models, this scenario is incomplete because it does not include an explanation of what precedes the smoothing phase. The anamorphic phase only occurs once and the universe expands forever in the future. Θ_{P1} is positive and monotonically decreasing throughout. This simple model is sufficient to explain smoothness, flatness, and the generation of a nearly scale-invariant spectrum of density perturbations without inflation and without producing a multiverse.




Figure 1. Sketch of the (Jordan-frame) potential, V_J (continuous black line), the gravitational coupling, f (dotted blue line) and the kinetic coupling, k (dashed green line) versus ϕ in the simple anamorphic scenario. During the smoothing phase, the anamorphic field ϕ moves uphill along the potential energy curve due to the wrong-sign kinetic term while f grows monotonically. The anamorphic phase ends when, on the way uphill, the field’s kinetic energy becomes comparable with its potential energy and $\epsilon \rightarrow 1$. The remaining kinetic energy of the field is sufficient to carry ϕ further uphill and over the top of the potential. On the way downhill, a transition to expansion occurs (Θ_m -bounce) when the kinetic coupling changes sign at $\phi_t \sim M_{\text{Pl}}^0$. After the Θ_m -bounce, ϕ reaches the minimum of the potential well and starts to oscillate rapidly, decaying and reheating the universe. Near the minimum, ϕ obtains a large mass and f and k become fixed so that the Einstein and Jordan frames become indistinguishable. The vacuum energy density at the minimum has a small positive value corresponding to the current dark energy density.

The second example we will describe in section 8.2 is a *cyclic anamorphic scenario* with repeated epochs controlled by a single scalar field that each include: (i) a dark energy dominated expanding phase (like the present universe) in which the field value is near zero; (ii) decay to a contracting phase during which both Θ_m and Θ_{Pl} undergo a reversal and become negative; (iii) a Θ_{Pl} -bounce in which Θ_{Pl} switches from negative to positive via a novel well-behaved NEC-violating phase; (iv) an anamorphic smoothing phase; (v) a Θ_m -bounce with reheating followed by a rapid return of the scalar field to its original value (near zero); (vi) normal hot big-bang expansion until the dark energy overtakes matter and radiation, which brings the cycle back to step (i). In principle, the cycles can continue forever following this sequence. The cyclic scenario has all the advantages of the simple model and also has the property that it can be made geodesically complete and thereby avoid initial condition problems.

8.1 Simple (one-time) anamorphosis

In this subsection, we describe a simple (non-cyclic) anamorphic scenario in which the universe begins in the anamorphic smoothing phase and subsequently reheats and transitions to a standard big-bang phase. Beginning in the anamorphic phase means that the condition in eq. (7.4) is satisfied initially.

In figure 1, we have drawn a typical anamorphic potential with gravitational and kinetic couplings. In figure 2, we have plotted the corresponding equation-of state parameter ϵ and




Figure 2. Plot of the equation-of-state parameter, ϵ (continuous black line in upper figure) and the frame-invariants, Θ_{PI} (dashed black line in lower figure) and Θ_m (continuous blue line in lower figure) as a function of $\phi \equiv \ln(A\chi)/A$ for the anamorphic Lagrangian (5.1) with $f(\chi) = \xi\chi^2/(1+\beta^2\chi^2)$, $k(\chi) = -1+2\gamma^2\chi^2/(1+\gamma^2\chi^2)$ and $V_J(\chi) = \lambda\chi^q \exp(-\alpha\chi)$ and the parameter values $\xi = 0.167$, $q = 3.99$, $\alpha = 1$, $\beta = 0.01$, and $\gamma = 0.05$. The dotted orange line in the lower figure is the (Jordan-frame) time-derivative of Θ_m . The dashed green line in the upper figure is the function defined in eq. (B.6), $3 + 2kf/f'^2$. When $3 + 2kf/f'^2 = \epsilon$ the transition from (ordinary) contraction to (decelerated) expansion occurs; ϕ_t ($\lesssim M_{PI}^0 \equiv 1$) is the field value at the transition. The y -axis of the lower figure has arbitrary units.

the frame-invariants, Θ_m and Θ_{PI} (all as a function of the anamorphic field ϕ) up to the transition to reheating and the hot big-bang phase. Although these are sketches, all the properties described by these figures and in the discussion below have been confirmed by explicitly solving specific examples, such as the one detailed in the caption to figure 2. The solutions were found numerically by solving the differential equation eq. (7.3) for ϵ as a function ϕ , and then substituting the solution into the following relations:

$$\Theta_{PI}(\phi) = \sqrt{\frac{V_J}{f^2(3-\epsilon)}}, \quad (8.1)$$

$$\Theta_m(\phi) = \Theta_{PI}(\phi) - \frac{1}{2} \frac{f_{,\phi}}{f} \frac{\dot{\phi}_J}{\sqrt{f}}, \quad (8.2)$$

$$\dot{\phi}_J(\phi) = \Theta_{PI} \sqrt{\frac{2\epsilon}{K}} f. \quad (8.3)$$

The first relation follows from rewriting the Friedmann equations re-expressed in terms of the invariant Θ_{P1} ; the second and third expressions follow from the definitions of Θ_m , Θ_{P1} and ϵ .

The key features of the simple scenario are as follows:

Anamorphic smoothing: the cosmological evolution in the simple scenario starts with the anamorphic smoothing phase at small field values (on the left hand side of the barrier in figure 1 where $\phi \ll \phi_t \lesssim M_{\text{P1}}^0 \equiv 1$) and the potential energy density being very low. In this phase, ϵ is less than one and slowly varying, as given by eq. (7.5). Also, $\Theta_m < 0$ (see lower panel of figure 2), corresponding to a universe that is contracting. Due to the wrong-sign of the kinetic term k the anamorphic field, ϕ moves uphill along the potential energy curve V_J . The gravitational coupling f grows monotonically and so the effective gravitational constant becomes weaker, contributing to the anamorphic smoothing process (recall eq. (4.5)). Θ_{P1} is positive and very slowly decreasing, ensuring that the curvature perturbations are (nearly) scale invariant with a small red tilt. The amount of anamorphic smoothing is given by the logarithmic change in α_{P1} , $\ln(\alpha_0/\alpha_{\text{P1}})$, where α_0 is the value of α_{P1} at the end of the smoothing phase, as in eq. (4.12). Quantum fluctuations generated during this period result in the scalar spectral tilt n_S and tensor-to-scalar ratio r computed in section 7.3, where the result (Case I or Case II) depends on the choice of parameters for V_J , f and k .

Exit and Transition: the anamorphic smoothing phase comes to an end when the kinetic energy becomes comparable to the field's potential energy, i.e., $\epsilon \simeq 1$. Although the anamorphic phase ends before ϕ reaches the maximum of the potential, ϕ continues to grow because the universe is still contracting (until $\phi = \phi_t$) and, hence, the kinetic energy of ϕ continues to blue shift. The value of ϵ increases over the next interval because of a combination of factors: the potential V_J has a maximum, as depicted in figure 1; the gravitational coupling f increases with ϕ ; and the kinetic coupling k is nearly constant ~ -1 (for $\phi < \phi_t$).

Θ_m reaches its most negative value and begins to increase after ϵ reaches $\epsilon \sim 1$ and the anamorphic phase ends. Over the next period, because $\dot{\Theta}_m < 0$, Θ_m becomes more and more negative, but now at a slower rate than before due to the fact that $\dot{\Theta}_m$ is increasing towards zero. Next, $\dot{\Theta}_m$ grows to the point that it reaches zero and, hence, $\Theta_m \ll 0$ reaches its most negative value. As $\dot{\Theta}_m$ passes through zero and becomes positive, the field is still climbing uphill but Θ_m is becoming less negative. When $\dot{\Theta}_m$ hits its maximum, ϕ reaches the top of the potential and begins to roll downhill towards the minimum of the potential V_J . At last, when the kinetic coupling, k , passes through zero and changes from negative to positive at $\phi = \phi_t$, Θ_m also passes through zero and becomes positive. This is the moment we call the Θ_m -bounce. (In the model proposed by Li [30], k remains negative throughout, including the hot expanding phase when the reduced Planck mass becomes fixed; an equivalent of the Θ_m -bounce is achieved by introducing tuned modifications of the effective potential which potentially induces an unintended inflationary phase that leads to a multiverse.)

Notably, the Θ_m -bounce can occur without violating the null-energy condition (NEC), as can be seen from eqs. (A.1)–(A.2). The Θ_m -bounce occurs at finite values of the effective Weyl-invariant scale factor α_m and does not require a Θ_{P1} -bounce. After the Θ_m -bounce, the field settles at the minimum of the potential where ϕ obtains a large mass; f and k become fixed such that $M_{\text{P1}}\Theta_m = M_{\text{P1}}\Theta_{\text{P1}} = H_E$ in agreement with Einstein gravity and standard big-bang evolution.

No multiverse: as discussed in section 6.2, the inflationary multiverse problem arises due to rare quantum fluctuations that kick the inflaton field in a direction that delays the end of the inflationary smoothing phase. Rare regions where this occurs continue to inflate and grow exponentially faster than regions that end inflation, so the rare regions soon occupy

the overwhelming majority of the volume of the universe. This process repeats *ad infinitum*, leading to runaway ‘eternal inflation’ and a multiverse. In ekpyrotic models, the problem is avoided because the analogous rare fluctuations extend the duration of the ekpyrotic smoothing, but regions in the smoothing phase are *contracting* in this case. The regions undergoing the rare fluctuations remain negligibly small compared to typical regions that complete the smoothing phase, bounce and expand, and so there is no runaway smoothing or multiverse.

In the anamorphic scenario, smoothing also occurs during a contracting phase, but the situation is more subtle than the ekpyrotic case. For some rare fluctuations, the only effect is to extend the contracting phase, just as in the ekpyrotic model, and the same argument ensures that these regions cause no problems. However, some rare fluctuations can cause the evolution of the scalar field to reverse direction (e.g., towards the left in figure 1) compared to the normal classical evolution in the anamorphic phase. In those regions, $\Theta_m = \Theta_{\text{PI}} - (f_{,\phi}/2f)(\dot{\phi}/M_{\text{Pl}})$ flips sign from negative to positive, corresponding to expansion. If the expansion were inflationary and sustained for many e -folds, this could potentially cause runaway and produce a multiverse. However, it is straightforward to show this does not occur. If a quantum fluctuation gives the scalar field a strong kick in the wrong direction and the kinetic energy density is large compared to the potential energy density, there is expansion but no inflation. Analytic estimates (and our numerical solutions) show that the expansion lasts for only a fraction of an e -fold before the field picks up enough speed for Θ_m to flip sign from positive to negative, leading to a long period of contraction. Because the expansion caused by the rare fluctuation does not even last for an e -fold, there is no chance of runaway inflation, and no multiverse problem arises.

8.2 Cyclic anamorphosis

In the preceding section we presented a one-time cosmological scenario where we started with the anamorphic smoothing phase, transitioned to standard big-bang evolution via a Θ_m -bounce and ended with the current phase of dark-energy domination. In this section, we briefly outline an example of how the anamorphic scenario can be made cyclic, i.e., how a dark-energy phase like the present epoch can be connected to an anamorphic smoothing phase that seeds initial conditions for the next cycle of evolution; for a detailed description see our analysis in [23]. A cyclic scenario has the advantage of avoiding initial condition problems related to assuming any sort of beginning in time. The key stages of the cyclic scenario are illustrated in figure 3:

Dark-energy domination (the present phase, $\phi = \phi_0 \simeq 10^{-16} M_{\text{Pl}}^0$): for the purposes of illustration, we shall take the anamorphic field to be the Higgs field, where we will assume that the current vacuum ($\phi = \phi_0$) is metastable and ϕ_0 has the empirically measured value. In the current cosmological phase, the anamorphic field is settled in a local, metastable vacuum state of the Higgs-like potential $V_J(\phi)$, separated by a barrier of $\sim (10^{10-12} \text{ GeV})^4$ from the true, negative energy-density vacuum. The potential energy density $V_J(\phi_0)$ is the dominant energy component and it has the observed value of the current vacuum or dark-energy density $\sim (10^{-12} \text{ GeV})^4$; the gravitational coupling is $f(\phi_0) \equiv 1$ (in agreement with tests of general relativity today) and the kinetic coupling is $k \equiv 1$ (in accordance with the standard Higgs model). Due to the positive vacuum energy density of the metastable vacuum, the cosmological background expands at an accelerated rate.

Contraction ($10^{-6} M_{\text{Pl}}^0 \lesssim \phi \lesssim \phi_{\text{NEC}}$): eventually, the metastable vacuum decays and ϕ rolls or tunnels to the negative part of the potential. As the field rolls downhill to more negative potential energy density, $\Theta_m = \Theta_{\text{PI}}$ both undergo a reversal from positive to negative




Figure 3. Cyclic anamorphosis. A sequence of expanded views of the (Jordan-frame) potential V_J (continuous blue line) versus the anamorphic field ϕ in the cyclic anamorphic scenario with the middle inset showing the energy barrier and the final inset showing the current metastable vacuum. Both the anamorphic field ϕ and its potential V_J are given in Planck units. We superposed curves to represent the gravitational coupling, f (dotted-dashed red line) and the kinetic coupling, k (dashed green line) that both are dimensionless functions of ϕ . For small field values ($\phi \ll \phi_{\text{ANA}}$) and, in particular, during the current cosmological epoch when the anamorphic field is trapped in the metastable vacuum, $f = 1$ and $k = 1$, in agreement with observations; the plots of f and k have been shifted apart for the purpose of illustration.

and the universe starts contracting. During the contracting phase that follows, the blue shifting kinetic energy density rapidly dominates the total energy density such that $\epsilon \gtrsim 3$. Rolling downhill, the anamorphic field keeps picking up kinetic energy and, due to its large kinetic energy density ($\epsilon \gg 3$) at the bottom of the potential, V_{min} , the field does not settle in the true, negative vacuum but continues to increase and roll uphill while the background keeps contracting.

NEC-violation and non-singular Θ_{PI} -bounce ($\phi_{\text{NEC}} \lesssim \phi \lesssim \phi_{\text{ANA}} \simeq M_{\text{Pl}}^0$): rolling uphill, the field's kinetic energy grows at a slower rate but continues to dominate ($\epsilon \sim 3$) for a substantial period after $V_J(\phi)$ changes from negative to positive. The depth of the potential $|V_{\text{min}}|$ can be chosen so that $\epsilon \sim 3$ when $\phi = \phi_{\text{NEC}}$ where $k(\phi)$ changes from positive to

negative, as illustrated in figure 3. The sign-switch of the kinetic coupling leads to a brief violation of the null-energy condition (NEC). As we show in [23], it is possible to achieve a ghost-free and stable NEC-violating phase by assuming that the effective Lagrangian for $\phi_{\text{NEC}} \lesssim \phi \lesssim \phi_{\text{ANA}}$ is Galileon-like [9–11, 14, 33] with higher order kinetic terms, $\sim (\partial\phi)^4, \sim \square\phi(\partial\phi)^2$. During NEC violation, $H_E = M_{\text{Pl}}\Theta_{\text{Pl}}$ begins less than zero, increases steadily and eventually hits zero and switches from negative to positive, resulting in a Θ_{Pl} -bounce. At the time of the Θ_{Pl} -bounce, the gravitational coupling f is beginning to increase, as depicted in figure 3. It is at this point in the evolution that Θ_m and Θ_{Pl} first become distinguishable. In particular, Θ_m remains finite and negative during and after the Θ_{Pl} -bounce, while Θ_{Pl} becomes positive. With increasing f , the quantity $(1/2)K(\phi)(\dot{\phi}/M_{\text{Pl}})^2$ as given in eq. (A.9) becomes positive and the NEC-violating phase ends. At this point, the couplings to higher order kinetic terms (functions of ϕ) can be chosen that the terms are negligible as ϕ increases further.

Anamorphic smoothing phase ($\phi_{\text{ANA}} < \phi$): the conditions are now precisely what is required for the anamorphic smoothing phase. That is, during the first three stages, the dark energy-dominated phase ends by tunneling to a state with negative potential energy density; a period of contraction begins that drives the field uphill in V_J through the brief period of NEC-violation that flips the sign of Θ_{Pl} to be positive and opposite that of Θ_m ; and the NEC-violating phase ends at approximately the value of $V_J(\phi)$ needed to generate energy-density perturbations of the right amplitude, as given by eq. (7.8). (Note that, the important quantity for obtaining the right perturbation amplitude is V_J/f^2 , which explains why it is possible in the simple scenario to have the anamorphic phase occur at small field values where V_J and f are small, whereas here the field value, $V_J(\phi)$ and f are all comparatively large.)

Due to the wrong-sign of the kinetic term k the anamorphic field ϕ continues to move uphill along the potential energy curve V_J . The gravitational coupling, f grows monotonically and so the effective gravitational constant becomes weaker, contributing to the anamorphic smoothing process. The Hubble-like parameters $\Theta_m < 0$ and $\Theta_{\text{Pl}} > 0$ have opposite sign, though, both Θ_m^2 and Θ_{Pl}^2 are very slowly shrinking and approaching zero during the anamorphic phase while the mass-variation index q as defined in eq. (4.7) remains nearly constant. Notably, $\dot{\Theta}_m/M_{\text{Pl}} > 0$ without violating the NEC. The very slow decrease of Θ_m^2 and Θ_{Pl}^2 ensures that the curvature perturbations are (nearly) scale-invariant with a small red tilt. The scalar and tensor spectra can be computed as described in section 7.3, where the result (Case I or Case II) depends on the choice of parameters for V_J , f and k .

Exit, Θ_m -bounce and hot big-bang evolution: the anamorphic smoothing phase ends when ϕ reaches infinity after finite time Δt_J and $\Theta_m < 0$ approaches zero. We assume a bounce occurs, analogous to that in ekpyrotic models, in which $\dot{\phi}$ reverses sign and gets a small kick [4]. Θ_m continues to increase, which means it switches from negative to positive resulting in a Θ_m -bounce. Consequently, the universe starts expanding. No Θ_{Pl} -bounce occurs, though. Rather Θ_{Pl} undergoes a reversal: it is positive before the Θ_m -bounce, reaches zero at the Θ_m -bounce and continues to decrease and becomes negative after the Θ_m -bounce.

On the way downhill, the gravitational coupling f is shrinking and the kinetic coupling k is growing, i.e., becoming less negative. The field eventually re-enters the range of NEC-violation encountered on the way uphill. This occurs, when $(f, \phi/f)^2$ has become small again relative to $|k/f|$. As before, $\dot{\Theta}_{\text{Pl}}/M_{\text{Pl}} > 0$ leads to a Θ_{Pl} -bounce and Θ_{Pl} becomes positive. No Θ_m -bounce occurs. As ϕ continues to decrease and \dot{f}/M_{Pl} vanishes, the invariants $\Theta_m, \Theta_{\text{Pl}}$ (now both positive) as well as the frames become indistinguishable. The field continues to

roll down the potential energy curve, reaches the negative minimum and continues uphill. The small extra kick generated at the Θ_m -bounce enables the field to cross the tiny barrier and settle in the metastable vacuum state. The Higgs field oscillates around its minimum converting its remaining kinetic energy to radiation, thereby reheating the universe. After 14 billion years of expansion and cooling, the universe reaches a condition like the present, dominated by the small, positive false vacuum energy density of the Higgs field, and the cycle returns to step 1. Note that, as in the ekpyrotic cyclic model [37], there is a long period of dark-energy domination such that fluctuations produced in previous cycles are redshifted to super-horizon wavelengths that are not visible in the next cycle.

9 Discussion

The goal of this paper has been to introduce a new approach for smoothing and flattening the early universe and generating a nearly scale-invariant spectrum of perturbations utilizing a single scalar field that couples non-minimally to gravity. Through the invariants Θ_m and Θ_{P1} , we have shown that we can classify different cosmological smoothing mechanisms and specify how the anamorphic approach differs from previously known inflationary and ekpyrotic approaches (which may be formulated with or without non-minimal coupling). It is clear from this classification that the anamorphic mechanism is a meld of the earlier ideas, aiming to employ the best features of both.

During the smoothing phase, the conditions on the equation of state ϵ and Θ_{P1} are reminiscent of inflation. Much of the intuition and experience gained in constructing inflationary models can be usefully applied in constructing models of the anamorphic smoothing phase. However, the differences are also important. The universe is physically contracting in the anamorphic phase, as quantified by Θ_m , which is key to avoiding the multiverse problem of inflationary models. Also, there are enormous differences in the conditions for embedding the smoothing phase in a complete cosmological model. Inflation relies on a big bang to precede inflation and leads to eternal inflation. To utilize the anamorphic mechanism, the big bang must be replaced by a Θ_m -bounce, and the anamorphic phase must precede that bounce.

In this respect, anamorphic cosmology is reminiscent of ekpyrotic cosmology. For example, as we illustrated in one of our constructions, replacing the bang with a bounce opens the possibility of embedding the anamorphic mechanism in a geodesically complete cyclic theory of the universe that resolves both the initial conditions and multiverse problems. (Notably, our construction includes a novel Galileon-like mechanism for producing a non-singular Θ_{P1} -bounce that evades gradient instabilities and ghosts, as detailed in ref. [23].)

The specific examples presented in this paper were constructed to illustrate certain design principles. In future papers, we will present examples that are more efficient and aesthetic. We will also show how to generalize the anamorphic mechanism to more general types of scalar-tensor and other modified gravity theories.

Acknowledgments

We thank D. Eisenstein, M. Johnson, E. Komatsu, S. Perlmutter and N. Turok for helpful discussions. This research was partially supported by the U.S. Department of Energy under grant number DE-FG02-91ER40671 (PJS).

A Cosmological background in Jordan and Einstein frames

We present the background equations of motion for the anamorphic action in eq. (5.1) in both Jordan and Einstein frames.

A.1 Jordan-frame representation

The Jordan frame with $M_{\text{Pl}}\Theta_m = H_J \neq M_{\text{Pl}}\Theta_{\text{Pl}}$ has the property that the dust particle mass m_J is independent of ϕ . Consequently, this is the frame in which physical rulers (set by the particle's Compton wavelength, say) have fixed length. Varying the action with respect to the metric and the field leads to the (Jordan-frame) equations of motion

$$3f(\phi)\left(H_J^2 + \frac{\kappa}{a_J^2}\right) - \frac{\sigma^2}{f(\phi)a_J^6} = \frac{1}{2}k(\phi)\dot{\phi}_J^2 + V_J(\phi) - 3H_J f_{,\phi}\dot{\phi}_J + \rho_m^J, \quad (\text{A.1})$$

$$f(\phi)\left(\dot{H}_J - \frac{\kappa}{a_J^2}\right) + \frac{\sigma^2}{f(\phi)a_J^6} = -\frac{1}{2}k(\phi)\dot{\phi}_J^2 + \frac{1}{2}H_J f_{,\phi}\dot{\phi}_J - \frac{1}{2}\left(f_{,\phi\phi}\dot{\phi}_J^2 + f_{,\phi}\ddot{\phi}_J\right) - \frac{1}{2}(\rho_m^J + p_m^J), \quad (\text{A.2})$$

$$k(\phi)(\ddot{\phi}_J + 3H_J\dot{\phi}_J) + V_{J,\phi} = 3f_{,\phi}\left(\dot{H}_J + 2H_J^2 + \frac{\kappa}{a_J^2} + \frac{\sigma^2}{3f^2 a_J^6}\right) - \frac{1}{2}k_{,\phi}\dot{\phi}_J^2, \quad (\text{A.3})$$

$$\frac{1}{2}\sum_i \dot{\beta}_i^2 = \frac{\sigma^2}{f^2(\phi)a_J^6}, \quad (\text{A.4})$$

where we define $f(\phi) \equiv (M_{\text{Pl}}(\phi)/M_{\text{Pl}}^0)^2$, where M_{Pl}^0 is the value of the reduced Planck mass today. The dot denotes differentiation with respect to the physical (Jordan frame) time, t_J , that runs from large negative to small negative values during the anamorphic phase; although the value of ϕ is frame invariant, time derivatives such as $\dot{\phi}_J$ are not so we add the J subscript to be clear that the derivative refers to Jordan frame time. ρ_m^J is the (Jordan-frame) matter energy density and p_m^J is the (Jordan-frame) matter pressure. The spatial curvature is $\kappa = (+1, 0, -1)$ and a homogeneous and anisotropic Kasner-like metric is assumed where

$$ds_J^2 = -dt_J^2 + a_J^2(t) \sum_i \exp(2\beta_i) dx_i^2 \quad \text{with} \quad \sum_i \beta_i(t) = 0. \quad (\text{A.5})$$

In the field equations, eq. (A.1)–(A.4), the anisotropy is parameterized by σ^2 where

$$\sigma^2 = \frac{1}{2} \sum_i c_i^2, \quad \beta_i = \frac{c_i}{f a_J^3}, \quad c_i \equiv \text{constant}. \quad (\text{A.6})$$

Smoothing and flattening the universe and generating nearly scale-invariant perturbations during the anamorphic phase and transitioning afterwards to a hot expanding phase imposes constraints on $f(\phi)$, $k(\phi)$ and $V_J(\phi)$. During the anamorphic phase, the invariant mass ratio m/M_{Pl} must shrink or, equivalently, the non-minimal coupling $f(\phi)$ must grow significantly to suppress the anisotropy. Then, during the hot expanding phase that follows, including the present epoch, $f(\phi)$ must approach unity so that $M_{\text{Pl}}^2 \rightarrow (M_{\text{Pl}}^0)^2 \equiv 1$. These conditions ensure that there are no observable deviations from Einstein gravity today or at observable red shifts.

In the hot expanding phase where $f(\phi)$ is constant, $k(\phi)$ must be positive (the standard sign in simple quantum field theories) so that the kinetic energy density is non-negative

and there is no ghost-field. However, during the anamorphic phase when $f(\phi)$ is growing, $k(\phi)$ must be negative in order to obtain the equation of state, ϵ , required for smoothing and flattening, as has been shown in section 5. Hence, the kinetic coupling $k(\phi)$ must change from negative during the anamorphic phase to positive during the subsequent hot expanding phase that follows. Because $k(\phi)$ varies from negative to positive during the scenario, it cannot be eliminated by a field re-definition in the Jordan-frame action. Smoothing also requires that the potential energy density $V_J(\phi)$ be positive and increasing during the anamorphic phase. This is achieved as a result of the negative (wrong-sign) kinetic coupling $k(\phi)$ during the anamorphic phase, which causes the solution for ϕ to run uphill along its potential $V_J(\phi)$. (By contrast, the analogous field during an ekpyrotic contracting phase has normal-sign kinetic energy density and runs downhill along a potential that is negative and decreasing.)

A.2 Einstein-frame representation

The Jordan frame is natural for describing the smoothing and flattening of the universe and the different stages of expansion and contraction before and after the smoothing phase because rulers have fixed lengths in this frame. However, for some purposes, it is useful to consider the action in the Einstein frame where $M_{\text{Pl}}\Theta_{\text{Pl}} = H_E \neq M_{\text{Pl}}\Theta_m$, especially when comparing gravitational effects, such as wavelengths of curvature perturbation modes compared to the Hubble radius. Under a Weyl transformation of the Jordan-frame metric, the Einstein-frame metric is

$$g_{\mu\nu}^E = f(\phi)g_{\mu\nu}^J. \quad (\text{A.7})$$

Introducing the field Φ given by

$$\frac{d\Phi}{d\phi} = \sqrt{K(\phi)}, \quad (\text{A.8})$$

where

$$K(\phi) \equiv \frac{\frac{3}{2}(f, \phi)^2 + k(\phi)f(\phi)}{f^2(\phi)}, \quad (\text{A.9})$$

the gravitational and field sector of the action can always be rewritten in terms of minimally-coupled Einstein gravity,

$$S_E = \int d^4x \sqrt{-g_E} \left(\frac{1}{2}R_E - \frac{1}{2}(\partial_\mu \Phi)^2 - V_E(\Phi) \right), \quad (\text{A.10})$$

where

$$V_E(\Phi) \equiv V_J(\phi)/f^2(\phi). \quad (\text{A.11})$$

During the anamorphic smoothing phase, both V_E and V_J are positive.

$K(\phi)$ must be positive definite and continuously differentiable in order for the field to have the right-sign kinetic term in the Einstein frame. Hence,

$$K(\phi) > 0 \quad (\text{A.12})$$

is the *no-ghost condition*. From the expression for $K(\phi)$ in eq. (A.9), we see that $k(\phi)$ can be negative and the theory can remain ghost-free if $(f, \phi)^2/f$ is sufficiently large.

A.3 Matter contribution and contraction

The dust contribution to the action in the two frames is such that the action of a single test particle is given as in eq. (2.1),

$$S_d = \int m_J ds_J = \int m_J / \sqrt{f(\phi)} ds_E. \quad (\text{A.13})$$

The dust-particle mass in Einstein-frame representation is given by

$$m_E = m_J / \sqrt{f(\phi)}. \quad (\text{A.14})$$

In the anamorphic models considered in this paper, m_J is ϕ -independent and, hence, m_E is ϕ -dependent. Because m_J is fixed independent of ϕ , having $H_J < 0$ (or, equivalently, $\Theta_m < 0$), as occurs during the anamorphic phase, means that the universe is contracting compared to physical rulers made from matter. However, as we have noted, the Hubble parameter in other frames may have a different sign. In particular, H_E (or, equivalently, Θ_{Pl}) is positive in the anamorphic phase.

B Conditions for successful anamorphosis

We analyze the conditions that $f(\phi) \equiv (M_{\text{Pl}}/M_{\text{Pl}}^0)^2$, $k(\phi)$ and $V_J(\phi)$ in the action (5.1) and that the equation of state $\epsilon(\phi)$ must satisfy in order to smooth and flatten the cosmological background and to have squeezed quantum fluctuations during the anamorphic contracting phase ($\Theta_m < 0 < \Theta_{\text{Pl}}$).

Constraint 1: contraction. The invariant anamorphic energy density is given by the sum of the kinetic and potential energy density of the anamorphic field ϕ and the invariant pressure is given by the difference:

$$\frac{\rho_A}{M_{\text{Pl}}^4} \equiv \frac{1}{2} K(\phi) \left(\frac{\dot{\phi}}{M_{\text{Pl}}^0 M_{\text{Pl}}} \right)^2 + \frac{V_J(\phi)/f^2(\phi)}{(M_{\text{Pl}}^0)^4}; \quad (\text{B.1})$$

$$\frac{p_A}{M_{\text{Pl}}^4} \equiv \frac{1}{2} K(\phi) \left(\frac{\dot{\phi}}{M_{\text{Pl}}^0 M_{\text{Pl}}} \right)^2 - \frac{V_J(\phi)/f^2(\phi)}{(M_{\text{Pl}}^0)^4}, \quad (\text{B.2})$$

where for pedagogical reasons we have kept factors of M_{Pl}^0 exposed. Since the energy density of the anamorphic scalar dominates the total energy density during the smoothing phase, the conventional equation-of-state parameter, $\epsilon \equiv (3/2)(1+w)$ with $w \equiv p/\rho$, can be re-expressed as

$$\epsilon \equiv \frac{1}{2} \frac{K(\phi)}{\Theta_{\text{Pl}}^2} \left(\frac{\dot{\phi}}{M_{\text{Pl}}^0 M_{\text{Pl}}} \right)^2, \quad (\text{B.3})$$

so that the anamorphic condition, $\Theta_m < 0$ (contraction relative to physical rulers) and $\Theta_{\text{Pl}} > 0$ (expansion relative to the Planck scale), becomes

$$\Theta_m = \Theta_{\text{Pl}} + M_{\text{Pl}}^{-1} \left(\frac{\dot{m}}{m} - \frac{\dot{M}_{\text{Pl}}}{M_{\text{Pl}}} \right) = \left(\sqrt{\frac{K(\phi)}{2\epsilon}} - \frac{1}{2} \frac{f_{,\phi}}{f} \right) \frac{\dot{\phi}_J}{M_{\text{Pl}}} < 0. \quad (\text{B.4})$$

Without loss of generality, we will consider the case when ϕ is increasing during the anamorphic phase, $\dot{\phi}/M_{\text{Pl}} > 0$. Since $K(\phi)$ and ϵ are positive definite, the only way Θ_m can be less than zero with $\dot{\phi}/M_{\text{Pl}} > 0$ is if

$$f_{,\phi} > 0. \quad (\text{B.5})$$

(The constraint is $f_{,\phi} < 0$ if $\dot{\phi}/M_{\text{Pl}} < 0$.) Then, substituting the definition of $K(\phi)$ from eq. (A.9) into eq. (B.4), the condition that $\Theta_m < 0$ reduces to

$$3 + 2k(\phi) \frac{f(\phi)}{f_{,\phi}^2} < \epsilon. \quad (\text{B.6})$$

It is straightforward to show that the constraint in eq. (B.6) is the same independent of the sign of $\dot{\phi}/M_{\text{Pl}}$.

Constraint 2: smoothing of the Cosmological Background. Smoothing and flattening during contraction requires that the scalar field energy density dominates over all other contributions for an extended period of 60 or more e -folds. We have shown that the necessary condition for smoothing and flattening is that $\epsilon_m > 1 - q$ or, equivalently, $\epsilon_{\text{Pl}} \equiv \epsilon < 1$ (see eqs. (4.10) and (4.16)). That is, smoothing and flattening imposes an upper-bound on ϵ which complements the lower bound constraint imposed by contraction, eq. (B.6).

Constraint 3: squeezing of quantum curvature perturbations. The squeezing constraint in eq. (4.11) can be rewritten as

$$\frac{d\alpha_{\text{Pl}}\Theta_{\text{Pl}}}{dt} M_{\text{Pl}}^{-1} = \alpha_{\text{Pl}}\Theta_{\text{Pl}}^2 \left(1 + \frac{d \ln \Theta_{\text{Pl}}}{d \ln \alpha_{\text{Pl}}} \right) > 0. \quad (\text{B.7})$$

Since $\epsilon \equiv d \ln \Theta_{\text{Pl}} / d \ln \alpha_{\text{Pl}}$, the squeezing constraint is equivalent to the smoothing condition $\epsilon < 1$.

Combining all three constraints with the no-ghost condition ($K > 0$), we have

$$0 < 3 + 2k(\phi) \frac{f(\phi)}{f_{,\phi}^2} < \epsilon < 1. \quad (\text{B.8})$$

References

- [1] PLANCK collaboration, P.A.R. Ade et al., *Planck 2015 results. XX. Constraints on inflation*, [arXiv:1502.02114](#) [INSPIRE].
- [2] A. Albrecht and P.J. Steinhardt, *Cosmology for Grand Unified Theories with Radiatively Induced Symmetry Breaking*, *Phys. Rev. Lett.* **48** (1982) 1220 [INSPIRE].
- [3] J.M. Bardeen, P.J. Steinhardt and M.S. Turner, *Spontaneous Creation of Almost Scale - Free Density Perturbations in an Inflationary Universe*, *Phys. Rev. D* **28** (1983) 679 [INSPIRE].
- [4] I. Bars, P.J. Steinhardt and N. Turok, *Cyclic Cosmology, Conformal Symmetry and the Metastability of the Higgs*, *Phys. Lett. B* **726** (2013) 50 [[arXiv:1307.8106](#)] [INSPIRE].
- [5] D. Baumann, P.J. Steinhardt, K. Takahashi and K. Ichiki, *Gravitational Wave Spectrum Induced by Primordial Scalar Perturbations*, *Phys. Rev. D* **76** (2007) 084019 [[hep-th/0703290](#)] [INSPIRE].
- [6] A. Borde, A.H. Guth and A. Vilenkin, *Inflationary space-times are incomplete in past directions*, *Phys. Rev. Lett.* **90** (2003) 151301 [[gr-qc/0110012](#)] [INSPIRE].
- [7] E.I. Buchbinder, J. Khoury and B.A. Ovrut, *New Ekpyrotic cosmology*, *Phys. Rev. D* **76** (2007) 123503 [[hep-th/0702154](#)] [INSPIRE].

- [8] E.I. Buchbinder, J. Khoury and B.A. Ovrut, *Non-Gaussianities in new ekpyrotic cosmology*, *Phys. Rev. Lett.* **100** (2008) 171302 [[arXiv:0710.5172](#)] [[INSPIRE](#)].
- [9] N. Chow and J. Khoury, *Galileon Cosmology*, *Phys. Rev. D* **80** (2009) 024037 [[arXiv:0905.1325](#)] [[INSPIRE](#)].
- [10] P. Creminelli, K. Hinterbichler, J. Khoury, A. Nicolis and E. Trincherini, *Subluminal Galilean Genesis*, *JHEP* **02** (2013) 006 [[arXiv:1209.3768](#)] [[INSPIRE](#)].
- [11] P. Creminelli, M.A. Luty, A. Nicolis and L. Senatore, *Starting the Universe: Stable Violation of the Null Energy Condition and Non-standard Cosmologies*, *JHEP* **12** (2006) 080 [[hep-th/0606090](#)] [[INSPIRE](#)].
- [12] P. Creminelli, A. Nicolis and M. Zaldarriaga, *Perturbations in bouncing cosmologies: Dynamical attractor versus scale invariance*, *Phys. Rev. D* **71** (2005) 063505 [[hep-th/0411270](#)] [[INSPIRE](#)].
- [13] G. Domènech and M. Sasaki, *Conformal Frame Dependence of Inflation*, *JCAP* **04** (2015) 022 [[arXiv:1501.07699](#)] [[INSPIRE](#)].
- [14] B. Elder, A. Joyce and J. Khoury, *From Satisfying to Violating the Null Energy Condition*, *Phys. Rev. D* **89** (2014) 044027 [[arXiv:1311.5889](#)] [[INSPIRE](#)].
- [15] F. Finelli, *Assisted contraction*, *Phys. Lett. B* **545** (2002) 1 [[hep-th/0206112](#)] [[INSPIRE](#)].
- [16] F. Finelli and R. Brandenberger, *On the generation of a scale invariant spectrum of adiabatic fluctuations in cosmological models with a contracting phase*, *Phys. Rev. D* **65** (2002) 103522 [[hep-th/0112249](#)] [[INSPIRE](#)].
- [17] M. Gasperini and G. Veneziano, *Inflation, deflation and frame independence in string cosmology*, *Mod. Phys. Lett. A* **8** (1993) 3701 [[hep-th/9309023](#)] [[INSPIRE](#)].
- [18] M. Gasperini and G. Veneziano, *Pre - big bang in string cosmology*, *Astropart. Phys.* **1** (1993) 317 [[hep-th/9211021](#)] [[INSPIRE](#)].
- [19] M. Gasperini and G. Veneziano, *Dilaton production in string cosmology*, *Phys. Rev. D* **50** (1994) 2519 [[gr-qc/9403031](#)] [[INSPIRE](#)].
- [20] G.W. Gibbons and N. Turok, *The Measure Problem in Cosmology*, *Phys. Rev. D* **77** (2008) 063516 [[hep-th/0609095](#)] [[INSPIRE](#)].
- [21] A.H. Guth, *The Inflationary Universe: A Possible Solution to the Horizon and Flatness Problems*, *Phys. Rev. D* **23** (1981) 347 [[INSPIRE](#)].
- [22] A.H. Guth, D.I. Kaiser and Y. Nomura, *Inflationary paradigm after Planck 2013*, *Phys. Lett. B* **733** (2014) 112 [[arXiv:1312.7619](#)] [[INSPIRE](#)].
- [23] Anna Ijjas and Paul J. Steinhardt, *Cyclic Anamorphic Cosmology*, in preparation.
- [24] A. Ijjas, P.J. Steinhardt and A. Loeb, *Scale-free primordial cosmology*, *Phys. Rev. D* **89** (2014) 023525 [[arXiv:1309.4480](#)] [[INSPIRE](#)].
- [25] J. Khoury, B.A. Ovrut, P.J. Steinhardt and N. Turok, *The Ekpyrotic universe: Colliding branes and the origin of the hot big bang*, *Phys. Rev. D* **64** (2001) 123522 [[hep-th/0103239](#)] [[INSPIRE](#)].
- [26] K. Koyama, S. Mizuno, F. Vernizzi and D. Wands, *Non-Gaussianities from ekpyrotic collapse with multiple fields*, *JCAP* **11** (2007) 024 [[arXiv:0708.4321](#)] [[INSPIRE](#)].
- [27] J.-L. Lehners, P. McFadden, N. Turok and P.J. Steinhardt, *Generating ekpyrotic curvature perturbations before the big bang*, *Phys. Rev. D* **76** (2007) 103501 [[hep-th/0702153](#)] [[INSPIRE](#)].
- [28] J.-L. Lehners and P.J. Steinhardt, *Intuitive understanding of non-Gaussianity in ekpyrotic and cyclic models*, *Phys. Rev. D* **78** (2008) 023506 [*Erratum ibid.* **D 79** (2009) 129902] [[arXiv:0804.1293](#)] [[INSPIRE](#)].

- [29] J.-L. Lehners and P.J. Steinhardt, *Non-Gaussian density fluctuations from entropically generated curvature perturbations in Ekpyrotic models*, *Phys. Rev. D* **77** (2008) 063533 [Erratum *ibid.* **D 79** (2009) 129903] [[arXiv:0712.3779](#)] [[INSPIRE](#)].
- [30] M. Li, *Generating scale-invariant tensor perturbations in the non-inflationary universe*, *Phys. Lett. B* **736** (2014) 488 [Erratum *ibid.* **B 747** (2015) 562] [[arXiv:1405.0211](#)] [[INSPIRE](#)].
- [31] A.D. Linde, *A New Inflationary Universe Scenario: A Possible Solution of the Horizon, Flatness, Homogeneity, Isotropy and Primordial Monopole Problems*, *Phys. Lett. B* **108** (1982) 389 [[INSPIRE](#)].
- [32] V.F. Mukhanov, H.A. Feldman and R.H. Brandenberger, *Theory of cosmological perturbations. Part 1. Classical perturbations. Part 2. Quantum theory of perturbations. Part 3. Extensions*, *Phys. Rept.* **215** (1992) 203 [[INSPIRE](#)].
- [33] A. Nicolis, R. Rattazzi and E. Trincherini, *The Galileon as a local modification of gravity*, *Phys. Rev. D* **79** (2009) 064036 [[arXiv:0811.2197](#)] [[INSPIRE](#)].
- [34] A. Notari and A. Riotto, *Isocurvature perturbations in the ekpyrotic universe*, *Nucl. Phys. B* **644** (2002) 371 [[hep-th/0205019](#)] [[INSPIRE](#)].
- [35] R. Penrose, *Difficulties with inflationary cosmology*, *Annals N. Y. Acad. Sci.* **571** (1989) 249 [[INSPIRE](#)].
- [36] Y.-S. Piao, *Conformally Dual to Inflation*, [arXiv:1112.3737](#) [[INSPIRE](#)].
- [37] P.J. Steinhardt and N. Turok, *A cyclic model of the universe*, *Science* **296** (2002) 1436 [[INSPIRE](#)].
- [38] P.J. Steinhardt, *Natural inflation*, in G.W. Gibbons, S. Hawking and S. Siklos eds., *The Very Early Universe*, Cambridge University Press, Cambridge, U.K. (1983), pp. 251–266.
- [39] A.J. Tolley and D.H. Wesley, *Scale-invariance in expanding and contracting universes from two-field models*, *JCAP* **05** (2007) 006 [[hep-th/0703101](#)] [[INSPIRE](#)].
- [40] G. Veneziano, *Scale factor duality for classical and quantum strings*, *Phys. Lett. B* **265** (1991) 287 [[INSPIRE](#)].
- [41] A. Vilenkin, *The Birth of Inflationary Universes*, *Phys. Rev. D* **27** (1983) 2848 [[INSPIRE](#)].
- [42] D. Wands, *Duality invariance of cosmological perturbation spectra*, *Phys. Rev. D* **60** (1999) 023507 [[gr-qc/9809062](#)] [[INSPIRE](#)].
- [43] L.-M. Wang, V.F. Mukhanov and P.J. Steinhardt, *On the problem of predicting inflationary perturbations*, *Phys. Lett. B* **414** (1997) 18 [[astro-ph/9709032](#)] [[INSPIRE](#)].
- [44] C. Wetterich, *Hot big bang or slow freeze?*, *Phys. Lett. B* **736** (2014) 506 [[arXiv:1401.5313](#)] [[INSPIRE](#)].